

The Arecibo Dual-Beam Survey: Arecibo¹ and VLA² Observations

Jessica L. Rosenberg & Stephen E. Schneider

Department of Astronomy, University of Massachusetts, Amherst, MA 01003

rosenber@umbriel.astro.umass.edu

schneide@messier.astro.umass.edu

ABSTRACT

The Arecibo Dual-Beam Survey is a “blind” 21 cm search for galaxies covering ~ 430 deg² of sky. We present the data from the detection survey as well as from the follow-up observations to confirm detections and improve positions and flux measurements. We find 265 galaxies, many of which are extremely low surface brightness. Some of these previously uncataloged galaxies lie within the zone of avoidance where they are obscured by the gas and dust in our Galaxy. 81 of these sources are not previously cataloged optically and there are 11 galaxies that have no associated optical counterpart or are only tentatively associated with faint wisps of nebulosity on the Digitized Sky Survey images. We discuss the properties of the survey and in particular we make direct determinations of the completeness and reliability of the sample. The behavior of the completeness and its dependencies is essential for determining the HI mass function. We leave the discussion of the mass function for a later paper, but do note that we find many low surface brightness galaxies and 7 sources with $M_{HI} < 10^8 M_{\odot}$.

1. INTRODUCTION

We present the Arecibo Dual-Beam Survey (ADBS), a “blind” 21cm survey for galaxies. “Blind” HI surveys provide a look at galaxy populations that is unbiased by stellar populations. Many previous searches have tried to determine whether there is a hidden population of HI-rich galaxies. They have looked in voids (Szomoru *et al.* 1996, Hulsbosch 1987, Krumm & Brosch 1984) in clusters (Banks *et al.* 1999, Szomoru *et al.* 1994, Weinberg *et al.* 1991), in groups (Kraan-Korteweg *et al.* 1999, Haynes & Roberts 1979, and Lo & Sargent 1979) and at random directions in the sky (Kilborn *et al.* 1999, Spitzak & Schneider 1998, Zwaan *et al.* 1997, Henning 1992, Fisher & Tully

¹The Arecibo Observatory is part of the National Astronomy and Ionosphere Center, which is operated by Cornell University under cooperative agreement with the National Science Foundation. in Puerto Rico.

²The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

1981). These surveys have not been able to convincingly determine the nature of the population of low HI-mass galaxies. Zwaan *et al.* (1997) and Spitzak & Schneider (1998) undertook “blind” HI surveys, similar to the one discussed here, but neither had enough low mass sources to make a strong claim about the shape of the mass function. Schneider *et al.* (1998) combined the two data sets and found evidence for a rise in the number of low mass sources, but the results are not definitive. The HIPASS survey, with its coverage of the entire Southern sky, should also contribute to our understanding of the HI mass function. The most recently published mass function contains 263 galaxies, comparable to the number in this survey (Kilborn *et al.* 1999) but only has 2 galaxies with $M_{HI} < 10^8 M_{\odot}$ (converting to $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$, which is used throughout this paper). Although the debate about the number of low HI mass sources persists, these galaxies certainly do exist. We detect 7 sources with HI masses $< 10^8 M_{\odot}$, almost as many as all of the previous “blind” HI surveys combined.

The ADBS is one of the largest and most sensitive surveys to date. Using the Arecibo telescope we were able to achieve an rms of 3-4 mJy at a resolution of 32 km s^{-1} in only 7 seconds and covered $\sim 430 \text{ deg}^2$. The ADBS covers substantially more volume than the deeper Spitzak & Schneider (1998) and the Zwaan *et al.* (1997) surveys, and is much more sensitive than the larger Parkes HIPASS survey (Kilborn *et al.* 1999). HIPASS will cover the entire southern sky when it is completed, but it has a sensitivity of only 13 mJy at a resolution of 13.2 km s^{-1} and has trouble with the confusion of multiple HI sources within its $14'$ beam. With the low sensitivity, the Parkes survey is only able to detect low mass sources at very low redshifts where distances are difficult to determine.

In this paper we will discuss the HI data and our efforts to quantify the completeness and reliability of this sample. This is the first survey to use synthetic sources to derive an empirical relation between the flux, line width, and completeness of the survey. This will allow us to determine an HI mass function even if we do not understand all of the circumstances under which sources are not detected. The HI mass function will be discussed in Rosenberg & Schneider (in preparation; hereafter paper II).

We also present the optical identifications of the ADBS sources from the Digitized Sky Survey (DSS) images. We identify 81 HI-rich galaxies that are not in optical catalogs. Out of these 81 sources, 11 are not identified because they are heavily obscured by the Galactic plane. An additional 11 sources are at high Galactic latitude where there is no clear optical identification. In most of these cases, there is a hint of nebulosity that might be associated, a possible background source, or a bright star, but no convincing association. Even if these optical counterparts are associated, they are extremely faint relative to the mass of HI present. These 11 sources represent the most extreme cases in the survey, but we detect many more sources that were previously uncataloged because they are very faint.

We describe the survey in §2 and the VLA and Arecibo follow-up in §3. The data and data tables are in §4 followed by a discussion of the completeness in §5 and the properties of the sample

in §6. The results of the survey are summarized in §7.

2. OBSERVATIONS

2.1. The Detection Survey

The goals of our survey are (1) to determine whether there are classes of galaxies that have been overlooked previously and (2) to help tie down the HI mass function with the larger number of sources in this sample. To achieve these goals a “blind” HI survey needs to detect galaxies down to the lowest possible HI fluxes. At least equally important is a detailed understanding of the survey limitations. The statistics of previous surveys have been insufficient to definitively establish the shape of the HI mass function at both the high and the low mass ends. To improve the statistics one needs to cover a large volume of space. For a fixed total observing time more angular coverage means less sensitivity but greater volume coverage for a given HI mass since $\text{area} \propto t^{-1}$ but $\text{depth} \propto t^{1/4}$ where t is the integration time per point. Therefore the effective volume covered in a fixed observing time $\propto t^{-1/4}$. This might suggest that integration times should be made as short as possible, except that as sensitivity drops, low mass sources are detectable at smaller redshifts where the distance uncertainty becomes more significant and confusion with Galactic HI becomes a problem.

In trying to detect sources there is also a trade off between completeness and reliability. As digging deeper into the noise improves completeness, but causes the selection of more and more false sources. Likewise, eliminating narrow features that appear to be interference will eliminate some narrow linewidth sources. A balance must be achieved between the number of sources that can be verified and going deep enough to detect some of the fainter sources. In the end, the importance is in quantifying these selection effects. We discuss our completeness and reliability in §5.

There are many factors that go into determining which telescope is most suitable for a given project including collecting area, beam size, and system temperature. The sensitivity is proportional to the collecting area and inversely proportional to the system temperature. The collecting area is also inversely proportional to the beam size, so the decrease in beam size for a large telescope increases the time required to cover a given area of the sky. If the system temperatures are the same, telescopes of different sizes can cover the same area with the same sensitivity in equal times, however if the beam size is too large then source confusion becomes a problem. Confusion may be resolved later, but then it may be difficult to determine whether the galaxies would have been detected independently. Fortunately, at the typical distance of sources detected in only 7 seconds of integration, Arecibo’s 3.3’ beamsizes is usually able to distinguish neighboring galaxies.

During these observations, Arecibo was being upgraded and had limited mobility. This was an advantage for this project because it made large amounts of telescope time available for driftscan searches. Observing in driftscan mode is an efficient method when the goal is to cover a large area

on the sky and Arecibo is sufficiently sensitive that useful depths can be reached in the time a source takes to drift through the telescope beam. The lack of tracking also meant that two feeds could be used simultaneously. We used the 21 cm and 22 cm circular polarization feeds, which have now been replaced by the Gregorian reflector. These were located on opposite corners of the carriage house and pointed 1.6° apart on the sky. Our ability to use two feeds doubled the area surveyed, although it did require coarsening the velocity resolution to twice that in the Slice Survey (Spitzak & Schneider 1998).

One of the major challenges of any single-dish HI survey is to distinguish between real sources and radio interference. We used three tactics to address this problem while also increasing the volume coverage and improving the signal-to-noise: (1) Comparing data from the 21 and 22 cm feeds taken simultaneously – the interference usually enters through far sidelobes and appears in both feeds while real sources appear in only one. (2) Comparing left and right circular polarizations in each feed – the average of the two polarizations provides improved signal-to-noise while the difference is an additional check for interference since it is often highly polarized. (3) Observing each declination strip twice, on separate days, providing confirmation of the source detections.

While the driftscan technique allowed us to cover a large volume of space rapidly, it also gave us uncertainties of up to $\sim 7'$ in the declination position because the sources could be in the main beam or first sidelobe (or rarely farther out if the source was very bright at 21 cm). The lack of declination information meant that the sources needed to be reobserved to determine their positions and fluxes. We subsequently followed-up the identifications at the Very Large Array (VLA) and at Arecibo (§3).

The Arecibo 305m telescope was dedicated to the ADBS for approximately 500 hours between December 1993 and February 1994. These data consist of $\sim 300,000$ spectra taken every 7 seconds and cover nearly 24 hours of time in each of 30 declination strips. The total sky coverage, shown in Figure 1, is approximately 430 deg^2 in the $3.3'$ main beam. In this figure each strip is surrounded by two thin lines showing the positions of the sidelobes. The thick lines denote the regions covered twice while the thin lines denote the regions of single coverage. The actual sky coverage is the convolution of the sky coverage in Figure 1 with the beam shape. Figure 2 shows this convolution for the beginning of a driftscan. The contours indicate the total integrated detection sensitivity at each position, normalized to unity along the beam’s center-line.

The velocity coverage for the survey was -654 to 7977 km s^{-1} . This coverage was achieved over 512 channels in each of two polarizations for each of two feeds, totalling 2048 channels. The channel spacing was 16.9 km s^{-1} and the resolution was 33.8 km s^{-1} after Hanning smoothing.

2.2. The ADBS Data Reduction

The data reduction for ADBS spectra was a several step process which was run as an external FORTRAN module to ANALYZ, Arecibo Observatory’s data reduction package. The data are

driftscans and the adjacent spectra behave as off scans for each spectrum. The resulting noise, after the data have been reduced, is 3-4 mJy. We review the basic steps taken in the data reduction here:

- (1) The total power in the spectra was adjusted according to the recorded power counters.
- (2) An average “off-source” spectrum was created from all of the spectra in a scan (usually 300) after eliminating any outliers—either due to interference signals in individual channels or continuum flux across all channels.
- (3) Each spectrum had the “off” subtracted and then was normalized by it: $(on - off)/off$.
- (4) Synthetic sources were added so that they would go through the flattening procedure. The synthetic sources are discussed further in §5.
- (5) The continuum emission was estimated from the total power counters for each spectrum and then subtracted using the bandpass shape of each feed. The bandpass is assumed to be Gaussian centered at 1408.5 and 1398.5 MHz respectively, with a half-power width of 52 MHz for both feeds. These values were tested and adjusted until subtraction gave a flat response.
- (6) After excluding channels with strong signals compared to surrounding channels, a linear fit was made to each spectrum and subtracted.
- (7) Two “off” scans were created from data on either side of each spectrum. These “offs” consisted of 19 spectra separated from the “on” spectrum by 6 spectra. If either of the “offs” produced an improvement they were used. The idea was to remove some of the longer time-scale problems (standing waves and interference), that are matched in nearby spectra without removing the source.
- (8) The spectra were Hanning smoothed.
- (9) At this point the baseline looked very flat in most cases but there were a few exceptions where there was still low level waviness to the baselines. A final polynomial fit was designed to eliminate these underlying variations while ignoring any regions where there were rapid variations. A fifth-order polynomial was fit to each spectrum excluding any regions where there was a change of 7 mJy ($\sim 2\sigma$) or greater over a range of $\sim 32\text{km s}^{-1}$. Tests showed that this yielded a highly constrained fit that ignored weak, narrow signals, although it could reduce the apparent flux from wide, weak signals, especially when they were near a bandpass edge.
- (10) Images were Hanning smoothed in right ascension.

The ADBS survey was analyzed both by-eye and using the SExtractor (Bertin & Arnouts 1996) computer algorithm. In order to facilitate both of these analyses, groups of 300 spectra were turned into two-dimensional grayscale FITS images. Figure 3 is an example of one of these images. The images are 601 rows high with 300 spectra from the 21 cm feed in the top half of the image and 300 22 cm feed spectra, taken at the same time as the first 300, in the bottom half. The topmost row of the image (row 601) gives the center velocity of each column or correlator segment. The

columns are broken into two sections; the first 512 columns represent the 512 correlator channels averaged over the 2 polarizations, the last 512 columns contain the absolute value of the difference between the polarizations used as an interference check. There are some extra columns between the sum and difference of the polarizations which contain information about the baseline fits for each spectrum, the strength of the continuum, and the spectrum coordinates so that the information is easily available while searching for galaxies.

There are a few features to note in Figure 3. The first is that the prominent black and white “candy cane” near the left edge of the image is the Milky Way. The Milky Way emission does not subtract completely from the difference between the two polarizations (located just to the right of the image center) probably because the response to such strong emission is not always linear. The vertical stripes through the averaged polarizations, which stand out better in the polarization difference, are due to continuous interference which is variable in strength. The leftmost interference stripe shows up in the polarization difference, but not in the averaged polarizations because it is weak and fairly constant in strength. Momentary interference appears as two blips, one in the 21cm feed and the other in the 22 cm feed. These blips are also apparent in the polarization difference image. Other images occasionally show a wavy background due to standing wave patterns set up in the telescope. In these regions, galaxy detection is difficult but the measured RMS noise is also correspondingly bigger. On rare occasions, there is such strong interference that galaxy detection is impossible. We have identified two galaxies in this image which are marked with arrows in Figure 3. These detections are average in brightness.

2.3. ADBS Source Identification

The two dimensional images (Figure 3) proved to be a good, albeit time consuming, way to look at these 300,000 spectra. The images allowed us to easily identify strong interference which showed up in both feeds and often persisted for several spectra or would appear at the same wavelength a little later. Without this larger scale view it also would have been easy to misidentify peaks in the standing wave patterns as sources. Each image was looked at in tandem with its pair from a different day. With two images, two feeds, and two polarizations at hand, the reliability of the galaxy identification was $\sim 58\%$ even for sources at a signal-to-noise of less than 7 (see §5 for full discussion of reliability). From the original ADBS data we selected 407 sources, 265 of which were later confirmed.

The identification of sources is a delicate balance between the need for clear-cut selection criteria and the desire for both completeness and reliability of the sources. Examining the data by-eye proved to be very effective in finding sources, but it can be difficult to quantify in terms of the selection criteria. We also used the SExtractor neural network detection algorithm (Bertin & Arnouts, 1996) to automate the detection process and make it easier to quantify. These results are discussed in §4. The first tests quickly proved that the automated procedure was not very good at providing a reliable data set given only a single image. Because of the noise characteristics

of the data, the program tended to find either myriad sources or none. The computer algorithm was much more effective when applied to the data for which there were two independent images. Cross-correlation of the detection positions dramatically reduced the number of spurious computer-identified sources. For those images where there was not a second observation, the human eye proved to be much more effective at selecting reliable source candidates.

3. FOLLOW-UP TO THE ADBS

3.1. The VLA Follow-Up

The first follow-up observations to the ADBS were made at 21 cm with the VLA in D-array made in January 1998. With an allotment of 24 hours, it was impossible to follow-up all of the ADBS sources so we opted to observe as many of the sources with 2 ADBS detections and velocities less than 3000 km s^{-1} as possible. We took 10 minute snapshots of 99 of the ADBS galaxies. There were 78 sources observed with velocities of less than 3000 km s^{-1} in this data set. Of the 99 sources observed, 90 were detected, proving this to be a highly reliable data set as expected for the sources with two ADBS detections.

The VLA D-array was used for these observations because the sensitivity of this configuration allowed us to detect our sources in a reasonable amount of time and provided good velocity resolution over a large field of view ($31'$ HPBW). The synthesized beam-width for this configuration was $44''$ and we observed at a velocity resolution of 20.5 km s^{-1} . With 10 minute snapshot integrations we were able to achieve an RMS noise of 1.35 mJy/beam at the primary beam center which is below the detection limits of the ADBS unless the source was extended over more than about 7 beams. The VLA also provides velocity field information which is unavailable with a single dish observation.

The VLA data reduction was performed using standard AIPS methods. The images were created and cleaned using IMAGR with natural weighting. The final images were then primary beam corrected (PBCOR). The resulting sources were fit using the Gaussian fitting routine JMFIT which determined the HI centers and sizes.

3.2. The Arecibo Follow-up

To refine the positions and fluxes for the remaining ADBS sources, they were observed with the newly refurbished Arecibo telescope in July 1998, May 1999, and August 1999. To determine the declinations and fluxes of our sources, we chose to track the telescope at the detection right ascension and drive the telescope $\pm 7.5'$ from the driftscan declination. We will refer to this observation method as “DECscan” mode. The telescope was driven in declination for a total of 3 minutes and spectra were dumped every 10 seconds providing 18 spectra separated by $50''$. The resulting RMS

sensitivity is 1.5 mJy. Figure 4a shows an example of the resulting data.

The flux was determined in each of the individual “DECscan” spectra and a Gaussian was fit to the resulting function of total fluxes versus spectrum position as in Figure 4b. The peak of the derived Gaussian gives the single-beam flux, and the full width at half-maximum of the fit is a measurement of the galaxy size. The fit center gives the improved declination position. The velocity width of the source was determined from a composite spectrum made up of the sum of the spectra in which the source was detectable.

For 10 of the survey sources (020045+2809, 024328+2035, 071028+2307, 084411+2208, 084504+0932, 113845+2008, 171634+2135, 180506+2308, 183229+2308, 231941+1011) the follow-up Arecibo “DECscans” confirmed the detections, but the association with an optical counterpart in the DSS images was in question. For these sources, an “RAscan” was performed, that was identical to the “DECscans” except that the telescope was fixed at the “DECscan” declination and was driven $\pm 7.5'$ in right ascension to improve the source coordinates. In most cases the tentative optical counterpart appeared to be confirmed. The source coordinates in Table 2 reflect the improved “RAscan” position.

A total of 378 sources were re-observed at Arecibo. Out of these 215 were detected, 31 had been previously detected at the VLA, and 14 sources were observed during two of the Arecibo observing runs. Multiple observations of the same sources were used to check the internal consistency of the data set (§6).

4. DATA

Table 1 contains a list of the detection statistics for the Arecibo Dual-Beam Survey. We identified 407 possible sources. The reliability of these sources was 78% when they were covered twice and 42% when there was only a single coverage. Of the 265 sources, 184 have previous optical identifications.

Table 2 contains all 265 of the sources that were re-detected on follow-up. The table contains the RMS of the detection scan or scans which is typically 3-4 mJy. The rest of the data in the table are the fluxes, line widths, and velocities from the Arecibo and VLA follow-up observations.

Table 3 is a summary of the ADBS catalog identifications and the optical data for these sources. This optical information is derived using SExtractor (Bertin & Arnouts, 1996) on the DSS images. The red plates of the POSS II (Second Palomar Optical Sky Survey) images were used when they were available, but for 48 sources the red POSS I (First Palomar Optical Sky Survey) plates were used as noted in the table.

SExtractor is a routine that first identifies sources in the image, extracts them, and determines their physical parameters. The routine determines the positions of the major and minor axes, and the total number of pixels exceeding 1σ . The dimensions, a and b, that SExtractor reports are the

RMS of the intensity along each axis. We determined the “size” of the galaxy from the number of pixels within the $1\text{-}\sigma$ level ellipse: $\text{Pixels} = \pi ab \cdot x^2$ where x is the factor to scale from RMS to the major or minor axis size. In most cases x was about 2.5. This sigma level seemed to best correlate with the visible extent of the galaxies, particularly the ones with extended low surface brightness regions.

Figure 5 shows a comparison between SExtractor sizes and UGC sizes. Overall, the SExtractor measurements agree fairly well with the UGC values. We suspect that most of the scatter is due to inconsistencies in the depth of the UGC measurement since they were done by-eye on the POSS I, where the resolution is lower and where the larger spacing between plates allowed for more variability due to vignetting. In a few of the most extreme cases, the claimed UGC dimensions appear to extend well beyond any visible emission from the galaxy.

Table 3 lists only those sources with optical identifications in the NASA/IPAC Extragalactic Database (NED³). Note that 9 of the identified sources come from low surface brightness or dwarf galaxy surveys like the LSBC (Schombert *et al.* 1992), ESDO (Eder *et al.* 1989), KK98 (Karachentseva & Karachentseva 1998), KDG (Karachentseva *et al.* 1996). There are also many larger low surface brightness sources in our lists that were identified in the UGC catalog (Nilson, 1973), but missed by more traditional magnitude-limited optical surveys. In addition, there are galaxies at low Galactic latitude and therefore hidden behind our Galaxy’s gas and dust that can not be detected optically. 30 of the 39 galaxies with visual extinctions > 1.0 are not in any of the optical catalogs.

5. COMPLETENESS AND RELIABILITY

Completeness is the percentage of galaxies detected in a given volume down to a defined sensitivity limit. It is important to understand the completeness of the survey as a function of its detection parameters because this relates the number of sources found to the number of sources actually present. Reliability is the percentage of sources initially selected that are subsequently confirmed. There is a balance between completeness and reliability. The more sources selected from the original data and the further into the noise one “digs,” the better the completeness. However, a larger fraction of the selected sources will be unreliable. Because of the telescope time required for confirmation, these two must be carefully balanced.

For the ADBS data we determine the reliability for both the by-eye and automated detection methods. The by-eye method proved to be much more reliable. Overall we confirm 78% of the possible signals with two original detections and 42% for those with only a single detection. The reliability is $>70\%$ for both single and double detections when the signal-to-noise is above 7 but it falls to 57% for double detections and 37% for single detections below this limit. While the

³The Database (NED) is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

reliability does decrease at lower signal-to-noise values, especially for sources with only a single detection, the statistics are still much better than with SExtractor. With SExtractor the reliability was only 12% for double detection sources and uncovered no genuine sources not already found by-eye.

In general, it is not possible to determine the completeness of a survey based on the detections. It is often assumed that the sensitivity of a survey is determined by the sources with the lowest signal-to-noise. Schneider *et al.* (1998) showed that $\mathcal{V}/\mathcal{V}_{max}$ tests indicate that this sort of simple sensitivity cutoff does not adequately describe survey completeness. To quantify the completeness of the ADBS we added synthetic sources spanning the full range of galaxy linewidths and with fluxes varying from undetectable to that of bright galaxies that we expected to detect easily. These sources were given realistic line profile shapes and were all added to the data prior to the reduction procedure.

We use the synthetic sources to determine completeness as a function of signal-to-noise and to compare the results of our by-eye and automated detection methods. Figure 6 shows the percentage completeness as a function of signal-to-noise for automated detection (solid line) and by-eye detection (dotted line). The figure demonstrates the similarity in detection ability of the two methods, but there is a small improvement using automated detection.

We use visual inspection in making our source lists because, though it slightly decreases the completeness, it greatly improves the reliability. We note that all of the by-eye detections that were confirmed were also detected by the automated algorithm. We also ran a test in which we reobserved 25 randomly-selected SExtractor detections. Out of these, 3 were also detected by-eye and were confirmed whereas none of additional 22 possible signals were confirmed in follow-up.

Two effects on completeness are apparent in these simulations. First, we confirm that wide-line sources are not detected at S/N levels quite as low as narrow-line sources when the noise is assumed to increase as $w^{0.5}$. By assuming the effective noise grows like $w^{0.75}$, which Schneider *et al.* (1998) adopted based on the $\mathcal{V}/\mathcal{V}_{max}$ test, a consistent shape for the completeness curve is found independent of profile width. Second, with the statistics of several hundred synthetic sources, it is possible to characterize the behavior of the roll-off in completeness. It is apparent that occasionally even high S/N sources are missed and some $< 5\sigma$ sources are detected. These results are consistent for the two detection methods and will be important for the determination of the mass function in paper II.

Figure 7 is another way of examining the completeness of the sample. The first panel shows the synthetic sources. The detected sources are shown as filled gray dots while those we did not detect are shown as open circles. The completeness line with a slope of 0.75 and a “S/N” of 5 (using the formalism of Schneider *et al.* 1998) is shown. The line represents a good fit to the lower limit of the detected points, but it **does not** represent a 100% completeness limit. The apparent difference in the completeness for narrower and wider sources is not a real one. In fact, the completeness above the “S/N” = 5 line is 78% for all sources and 80% for sources with widths $< 300 \text{ km s}^{-1}$.

The second panel in Figure 7 shows fluxes and line widths of confirmed real sources in the ADBS (gray dots). The data represented here are the confirmed values for the widths and fluxes adjusted for frequency response and the offset of the source from beam center. We use these confirmed sources since they are of higher signal-to-noise and therefore give a better indication of the true flux. We have eliminated all sources with offsets greater than $2'$ from this plot because of the uncertainties in the measured flux when sources are beyond the half-power beamwidth. These uncertainties arise from the effect of small errors in the pointing and a lack of complete information about the HI extent.

Figure 8 shows the percentage completeness for the synthetic sources as a function of velocity. Notice the spikes in incompleteness near either end and in the center of the velocity range. These correspond to the ends of the bandpass segments. The baseline fitting at the ends of the segments is particularly prone to problems with source subtraction because there is not baseline on either side of the feature to distinguish a source from a baseline “wobble.” There is the additional problem for real sources in these regions that the sensitivity is falling off rapidly at the bandpass edges. If we eliminate sources with velocities $< 300 \text{ km s}^{-1}$ (to deal with confusion with the Milky Way too) and between 3100 and 3600 km s^{-1} the data are 82% complete above our “ 5σ completeness” limit.

The ideal check of completeness is to have a much deeper survey over the same area of sky, with the same instrument, with which to compare your results. Zwaan *et al.* (1997) provide us with just such a comparison sample for a small section of our survey. Their sample consists of ~ 10 hour driftscans at 2 declinations covered repeatedly to achieve about 5 times the sensitivity of our survey (RMS noise = 0.75 mJy in 16 km s^{-1} channels). One of these strips, at $\sim 23^\circ$, overlaps one of ours within $20''$. Because their data are much more sensitive, we can test our completeness although small number statistics make it impossible to use these results for a detailed study of sensitivity roll-off.

The triangles in Figure 7 are the Zwaan *et al.* sources; the filled triangles are the sources we detect, the open triangles are the ones we miss. We have detected all of the Zwaan *et al.* sources above the “S/N” = 5 line. There are two sources that we pick up as a single source. They are plotted with the rest of the sources and one falls with the detections, so we probably would have detected it on its own, while we probably would have missed the fainter one.

6. PROPERTIES OF THE SAMPLE

6.1. Position Determination

Figures 9a and 9b provide an evaluation of the positional accuracy of the ADBS, VLA, and “DECscan” measurements. We compare positions to the optical centers when the optical association was clear or to the VLA positions when the association was not. The optical centers were determined from the DSS images. Figure 9a shows the uncertainty in the RA measurements. The

open histogram is the offset between the detection positions and either the VLA or the optical positions. These offsets have a standard deviation of $57''$, about half of the $105''$ separation between spectra in the detection survey. The shaded histogram is the offset between the VLA positions and the optical positions, these offsets have a standard deviation of $31''$. This standard deviation is larger than expected from the measurement uncertainties of the optical and VLA positions and may be partially due to asymmetries in the HI distribution with respect to the optical emission.

Figure 9b shows the uncertainty in the original declination positions also using either the VLA or optical counterpart positions. The open histogram is the offset between the detection declinations and either the VLA or the optical positions. The standard deviation in this distribution is $2.1'$ which is about two-thirds of the width of the main beam, but there are several sources that are detected in the sidelobes up to $7'$ from the VLA or optical centers. The sources with the biggest offsets are all large galaxies that we catch the edge of. The shaded histogram shows the offsets between the Arecibo “DECscan” positions and the VLA or optical positions. The standard deviation in these offsets is $29''$, similar to the VLA positional uncertainties (which are not shown, but are the same as for the right ascension). Figures 9a and 9b demonstrate that the final positions have uncertainties of $\sim 30''$ for the VLA and “DECscan” coordinates, but $2.1'$ in right ascensions for sources that were not re-observed at the VLA.

Figure 10 is an example of why our detection method does not always determine the best position for an object. The galaxy in this figure was detected in a slice that passed to its north. Because of the galaxy’s position angle, the detection position was too far to the east. The “DECscan” improved the measurement, but it was still affected by the galaxy’s position angle. For the larger galaxies, these positioning problems can cause substantial errors.

6.2. Flux Determination

The flux measurement for a source will be underestimated if it is offset from the center of the beam. As we have shown in §6.1, our initial ADBS positions are up to $7'$ from the galaxy center. Our follow-up observations have much higher S/N than the detection scans because the integration time is longer and because the sources are better centered. We have used several methods to evaluate the accuracy and the internal consistency of the follow-up flux measurements. The crosses in Figure 11a show the fluxes of Arecibo sources measured in both July 1998 and May 1999, showing the consistency of the measurements on 2 independent dates. The filled circles demonstrate the consistency between an on-off and the original “DECscan” value. To further test the accuracy of the flux determinations, we made “DECscan” observations of several UGC galaxies for which there are high quality, single-beam HI data in the literature. Figure 11b shows the relationship between our flux determination and the literature values for these UGC galaxies. Overall, our fluxes agree well with the literature values although a few of the literature measurements are somewhat high. We reobserved three of the most discrepant sources with on-off measurements and found values in better agreement with our “DECscan” fluxes suggesting that our fluxes are not underestimated.

6.3. Spatial Distribution

The ADBS survey covers velocities from $-654 < v < 7977 \text{ km s}^{-1}$. Figure 12 shows the distribution of HI sources with redshift. No sources were found at negative velocities in the sample, and only three sources were detected with $v > 7500 \text{ km s}^{-1}$. At low velocities, we avoided selecting high velocity clouds that were highly extended in right ascension. We found a few candidate signals in this velocity range, but none were confirmed. Figure 12 shows that the distribution of galaxies in velocity space is fairly flat; this is probably due to surveying many different environments over a large area on the sky which tends to smear the signatures of large scale structure.

The redshift-position distributions of the galaxies are shown in Figures 13 a and b. We have broken the sample into a low declination region and a high declination region and plot the redshift versus RA. The stars in the plot represent ADBS sources that are overlayed on gray dots representing the RC3 (de Vaucouleurs *et al.* 1991) sources. It is evident from these figures that, at least qualitatively, the ADBS sources follow the structures defined by the RC3 galaxies. This is consistent with the findings of Spitzak & Schneider (1998) and Zwaan *et al.* (1997) for their “blind” HI samples.

6.4. Optical Counterparts

The DSS provides us with a look at the optical nature of our survey sources. The POSS I and POSS II sizes are comparable, although the relationship has a lot of scatter. Sources with only POSS I images are footnoted in Table 3. We find a wide range of optical properties for the galaxies detected in this survey. The galaxy types range from average spirals with angular diameters from $15.5'$ (NGC 3628) to compact galaxies only a few arcseconds across to galaxies that appear to be extremely faint patches of nebulosity. Figure 14 shows a histogram of the angular sizes of the survey sources. Some sources, like those with just a bit of wispy nebulosity and low Galactic latitude objects, were not measurable optically so they are not included. While the mean size is $\sim 100''$, the majority of the sources are smaller than $70''$.

Table 4 lists all of the sources for which no optical counterpart was firmly identified. There are 11 galaxies with $A_v > 2 \text{ mag}$ that are not discernible on the POSS plates. All but one of these are in the Zone of Avoidance (ZOA) hidden by the gas, dust, and stars of our own galaxy. The one exception is an unusual case of a source at -34° latitude which is behind a very small dense clump of gas identified in the Schlegel *et al.* survey (1998) to have an $A_v > 6 \text{ mag}$. There are also 11 sources in the list which are not in fields of high extinction where an optical identification is not obvious. Figure 15 shows the DSS images for 4 of these extremely faint or optically ambivalent, in the case of 142335+2131, galaxies. We note that these 11 sources do not include several other very low surface brightness counterparts where we feel the match is unambiguous.

6.5. Masses

A thorough analysis of the HI and dynamic masses for the ADBS galaxies is left for paper II, but Figure 16 shows the distribution of HI masses to demonstrate the range of source types we detect. The masses are determined using the standard conversion from flux:

$$M_{HI} = 2.36 \times 10^5 \cdot D^2 \cdot \int S dv \quad (1)$$

The distance used to determine the HI mass is corrected by the potential flow algorithm, POTENT (Bertschinger *et al.* 1990), or V_0 for galaxies where the correction was undefined. The sample contains sources ranging in mass from 2.3×10^7 to $1.5 \times 10^{10} M_\odot$. There are 7 galaxies with masses $< 10^8 M_\odot$. This is relative to 4 sources in Spitzak & Schneider (1998) and 2 for Zwaan *et al.* (1997; assuming $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

7. SUMMARY

The size of HI identified samples is growing with the contributions of Zwaan *et al.* (1997), Spitzak & Schneider (1998), HIPASS (Kilborn *et al.* 1999) and this survey. This driftscan survey has contributed a sample of 265 sources to the endeavor, 81 of which were not previously cataloged. All of these sources have been confirmed and have improved positions and fluxes based on follow-up VLA and upgraded-Arecibo measurements.

More important than numbers detected first optically or in HI is a thorough understanding of completeness. To make an accurate assessment of the completeness of our sample we have used synthetic sources which were incorporated into the data prior to the data reduction. We are not 100% complete at any signal-to-noise level, but we have developed a clear picture of the sensitivity roll-off.

We have evaluated the positional accuracy of the survey and find that the original detection survey has a positional accuracy of $\sim 1'$ in RA and $\sim 2'$ in DEC. The follow-up VLA and Arecibo scanning measurements improve the positions to an accuracy of $\sim 30''$. These positions have allowed us to measure the probable optical counterparts in all but 22 cases, half of which are in the Galactic Plane.

Using the DSS we have examined the optical nature of the HI sources detected in this survey and find a wide range of optical properties for the galaxies. There are Messier objects, there are faint wisps of nebulosity, and there are 11 galaxies completely enshrouded in the dust of the Milky Way so as to be indiscernible on the DSS images. A second set of 11 sources have no clear optical counterpart or wisps of nebulosity that may be associated.

The flux measurements for these galaxies are repeatable and are in good agreement with values

from the literature. The detection fluxes, and in a few cases the confirmation fluxes, can be affected by the offset of the galaxy from beam center particularly for large galaxies. There does not, however, appear to be any discrepancy between fluxes determined from our special scanning technique and standard on/off measurements.

From a qualitative look at the spatial distribution of our sample of galaxies relative to the distribution of sources in the RC3 (de Vaucouleurs *et al.* 1991), it is evident that the HI-selected sources follow the structures defined by the RC3 sample. This is consistent with the findings of Spitzak & Schneider (1998) and Zwaan *et al.* (1997) for their “blind” HI surveys.

The masses of galaxies in this survey range from $2 \times 10^7 M_\odot$ to $1 \times 10^{10} M_\odot$. We find 7 galaxies with masses $< 10^8 M_\odot$ – almost as many as all of the previous “blind” HI surveys combined. This data set, in conjunction with the samples of Spitzak & Schneider (1998) and Zwaan *et al.* (1997), should help constrain the faint end of the HI mass function. The larger number statistics and use of synthetic sources to quantify our detection limits should prove extremely powerful in applying this data to determining the HI mass function. Paper II will investigate the mass function as determined by these galaxies in detail.

We would like to thank the staffs at Arecibo and the VLA for their assistance with the observations. We would also like to thank Frank Briggs and Ertugrul Sorar for helpful discussions about this work. This work was supported in part by NSF Presidential Young Investigator award AST-9158096, and 2 Sigma-Xi Grants-in-Aid of Research.

The Digitized Sky Surveys were produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166. The images of these surveys are based on photographic data obtained using the Oschin Schmidt Telescope on Palomar Mountain and the UK Schmidt Telescope. The plates were processed into the present compressed digital form with the permission of these institutions.

The National Geographic Society - Palomar Observatory Sky Atlas (POSS-I) was made by the California Institute of Technology with grants from the National Geographic Society.

The Second Palomar Observatory Sky Survey (POSS-II) was made by the California Institute of Technology with funds from the National Science Foundation, the National Geographic Society, the Sloan Foundation, the Samuel Oschin Foundation, and the Eastman Kodak Corporation.

The Oschin Schmidt Telescope is operated by the California Institute of Technology and Palomar Observatory.

The UK Schmidt Telescope was operated by the Royal Observatory Edinburgh, with funding from the UK Science and Engineering Research Council (later the UK Particle Physics and Astronomy Research Council), until 1988 June, and thereafter by the Anglo-Australian Observatory. The blue plates of the southern Sky Atlas and its Equatorial Extension (together known as the SERC-J), as well as the Equatorial Red (ER), and the Second Epoch [red] Survey (SES) were all

taken with the UK Schmidt.

REFERENCES

- Banks, G. D., Disney, M. J., Knezek, P. M., Jerjen, H., Barnes, D. G., Bhatal, R., de Blok, W. J. G., Boyce, P. J., Ekers, R. D., Freeman, K. C., Gibson, B. K., Henning, P. A., Kilborn, V., Kolibalski, B., Kraan-Korteweg, R. C., Malin, D. F., Minchin, R. F., Mould, J. R., Oosterloo, T., Price, R. M., Putman, M. E., Ryder, D., Sadler, E. M., Staveland-Smith, L., Stewart, I., Stootman, F., Vaile, R. A., Webster, R. L. & Wright, A. E., 1999 astro-ph/9906146.
- Bertin, E. & Arnouts, S. 1996, A&A, 117, 393.
- Bertschinger, E., Dekel, A., Faber, S. M., Dressler, A., & Burstein, D. 1990, ApJ, 364, 370.
- de Vaucouleurs G., de Vaucouleurs A., Corwin H.G., Buta R.J., Paturel G., & Fouque P. 1991, Third Reference Catalogue of Bright Galaxies (RC3) (New York: Springer-Verlag).
- Eder, J. A., Schombert, J. M., Dekel, A., Oemler, A. 1989, ApJ, 340, 29.
- Haynes, M. P. & Roberts, M. S. 1979, ApJ, 227, 767.
- Helou, G., Hoffman, G.L., Salpeter, E.E. 1984, ApJS, 55, 433.
- Henning, P. A. 1992, ApJS, 78, 365.
- Hulsbosch, A. N. M. 1987, A&AS, 69, 439.
- Karachentseva, V. E. & Karachentseva, I. D. 1998, A&AS, 127, 409.
- Karachentseva, V. E., Prugniel, P., Vennik, J., Richter, G. M., Thuan, T. X., & Martin, J. M. 1996, A&AS, 117, 343.
- Kilborn, V., Webster, R. L., & Staveland-Smith, L. 1999, PASA, 16, 1.
- Kraan-Korteweg, R. C., van Driel, W., Briggs, F., Binggeli, B., & Mostefaoui, T. I. 1999, A&AS, 135, 255.
- Krumm & Brosch 1984. AJ, 89, 1461.
- Lo, K. Y. & Sargent, W. L. W. 1979, ApJ, 227, 756.
- Nilson, P. 1973, Uppsala General Catalog of Galaxies (UGC) (Acta Universitatis Upsaliensis. Nova Acta Regiae Societatis Scientiarum Upsaliensis - Uppsala Astronomiska Observatoriums Annaler, Uppsala: Astronomiska Observatorium).
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525.

- Fisher, J. R. & Tully, R. B. 1981, *ApJ*, 243, L23.
- Schneider, S. E., Spitzak, J. G., Rosenberg, J. L. 1998, *ApJ*, 507, 9.
- Schneider, S. E., Thuan, T. X., Magri, C., & Wadiak, J. E. 1990, *ApJS*, 72, 245.
- Schombert, J. M., Bothun, G. D., Schneider, S. E., & Mcgaugh, S. S. 1992, *AJ*, 103, 1107.
- Spitzak, J. G. & Schneider, S. E. 1998, *ApJS*, 119, 159.
- Szomoru, A., Guhathakurta, P., Van Gorkom, J. H., Knapen, J. H., Weinberg, D. H., & Fruchter, A. S. 1994, *AJ*, 108, 491.
- Szomoru, A., Van Gorkom, J. H., & Gregg, M. D. 1996, *AJ*, 111, 2141.
- Warmels, R. H. 1986, PhD Thesis, University of Groningen.
- Weinberg, D. H., Szomoru, A., Guhathakurta, P., & Van Gorkom, J. H. 1991, *ApJ*, 372, 13.
- Zwaan, M., Briggs, F. H., Sprayberry, D. & Sorar, E. 1997, *ApJ*, 490, 173.

Fig. 1.— The sky coverage of the ADBS survey. The thick lines show the areas where the sky was covered twice, the thin lines show where it was covered only once. The narrow lines on either side of each coverage strip show the positions of the sidelobes.

Fig. 2.— A cross-section of the effective convolved sensitivity of the Arecibo beam as it moves across the sky during a driftscan.

Fig. 3.— An example of one of the FITS images created by stacking 300 spectra. The x-axis represents the velocity, or distance, dimension while the y-axis represents RA. The image is split vertically into the two feeds which are actually spectra of different regions of the sky observed simultaneously. Along the horizontal, there is a pattern in the center of the image which is where additional information about the spectra has been stored. To the left of that, the data displayed are the average of the two polarizations, while to the right the image is the absolute value of the polarization difference.

Fig. 4.— (a) Eight spectra (out of 18 total) in which a source was detected in a “DECscan.” The y-axis is a measurement of the flux of the source but each spectrum is offset for display purposes. (b) The flux in each of the 18 spectra. This function is fit with a gaussian to determine the flux and declination of the source, as well as a rough measure of the HI extent.

Fig. 5.— Comparison of SExtractor sizes and UGC sizes. The most errant points are generally low surface brightness galaxies for which SExtractor terminates too early, or for which the extent of the galaxy on the POSS I images, upon which the UGC was based, was unclear.

Fig. 6.— Completeness of the ADBS detection as determined by-eye (dotted line) and by computer algorithm (solid line) using synthetic sources inserted in the data.

Fig. 7.— The left-hand panel shows the relationship between flux and line width for the synthetic sources inserted into our data. The gray, filled circles are the sources we detect. The open circles are the sources we miss. The solid line represents the completeness limit of the survey as defined in Schneider *et al.* (1998) with a S/N constant of 5 and a slope of 0.75. The figure shows that even above the “completeness” limit we miss 20% of the sources. The second panel shows the same thing for real data. The filled dots are the sources detected in our survey. The filled triangles are the sources from Zwaan *et al.* (1997) that we also detect. The open triangles are the Zwaan *et al.* sources that we miss. The gray triangles are 2 galaxies detected as one in both our survey and Zwaan *et al.*. We probably would detect the brighter source by itself, but not the fainter one.

Fig. 8.— A histogram showing the percentage of sources above our “completeness” limit that are missed in each velocity bin.

Fig. 9.— (a) The open histogram in shows the difference between the RA detection positions and the VLA or optical centers. The filled histogram shows the difference in RA between the VLA and the optical centers. (b) The open histogram shows DEC difference between the detection positions and the VLA or optical centers. The filled histogram shows the difference between the “DECscan” positions and the VLA or optical centers.

Fig. 10.— A demonstration of what can happen when determining the position of a galaxy which is not aligned with a cardinal direction. The original driftscan determined the incorrect right ascension and our “DECscans” do not provide any additional information about the East-West location of the galaxy.

Fig. 11.— (a) A comparison between the Arecibo fluxes measured during each of the two Arecibo follow up runs is shown by the crosses. The filled circles show a comparison between the “DECscan” flux and an on-off measurement of the same source.(b) A comparison between the flux values measured in this survey and the literature values for UGC sources that are cataloged and have high quality Arecibo flux measurements in the literature. The symbols in the plot refer to papers as follows: $\times$ Haynes & Giovanelli 1984, $\circ$ Schneider *et al.* 1990, $\triangle$ Bica & Giovanelli 1986, $*$ Giovanelli *et al.* 1986, $\bullet$ Freudling *et al.* 1988, $\blacksquare$ Lewis 1987, and $\square$ Giovanelli & Haynes 1989.

Fig. 12.— Histogram of the heliocentric velocities for the galaxies detected in the ADBS survey.

Fig. 13.— The redshift distribution of the ADBS galaxies, black stars, relative to RC3 sources, gray dots, out to $cz = 8000 \text{ km s}^{-1}$. (a) The declination range 8° to 18° . (b) The declination range 18° to 28° .

Fig. 14.— Histogram of the SExtractor determined optical diameters from the DSS images.

Fig. 15.— The DSS images of 4 galaxies with faint, questionable optical counterparts.

Fig. 16.— Histogram of the HI masses of the ADBS sources.

Table 1. Detection Statistics

	N	1 observation	2 observations
Total Linear Coverage	...	132.3 hrs	399.3 hrs
Suspected Sources	407	146	261
Not Confirmed	142	85	57
Confirmed	265	61	204
Cataloged	184	50	134
Not Cataloged	81	11	70

Table 2. Galaxy Detection Data


Name	N_{obs} ¹	RA (2000)	DEC (2000)	RMS ² mJy	RMS ² Jy km s ⁻¹	A Flx ³ km s ⁻¹	A ΔV_{20} ⁴ km s ⁻¹	A ΔV_{50} ⁵ km s ⁻¹	A Vel ⁶ Jy km s ⁻¹	V Flx ⁷ km s ⁻¹	V ΔV_{50} ⁸ km s ⁻¹	V Vel ⁹ km s ⁻¹
000330+2312	2	00 03 30.5	23 12 03.6	3.41	3.39	4.663	363.355	338.732	7313.468	...	...	...
000407+2234	2	00 04 07.0	22 34 55.2	3.49	3.51	2.858	226.510	219.58	4475.981	...	...	...
000623+2347	2	00 06 23.0	23 47 31.2	3.57	3.62	2.618	337.622	296.021	4682.532	...	...	...
000900+2348	2	00 09 00.5	23 48 57.6	3.53	3.67	2.233	279.771	237.790	4492.880	...	...	...
001622+2237	2	00 16 22.6	22 37 04.8	3.54	3.51	3.416	325.708	249.79	5784.375	3.322	252.400	5819.007
002249+2310	2	00 22 49.2	23 10 19.2	3.42	3.45	1.179	48.448	34.961	4525.693	...	...	...
002526+2136	2	00 25 26.4	21 36 14.4	3.50	3.56	1.582	142.456	116.464	4631.979	3.025	169.800	4592.007
003426+2436	2	00 34 26.6	24 36 03.6	3.44	3.52	3.810	164.503	146.775	5341.837	...	...	...
003751+0838	2	00 37 51.4	08 38 31.2	3.64	3.68	5.826	394.050	355.016	5278.171	...	...	...
003811+2523	2	00 38 11.0	25 23 45.6	3.46	3.43	1.372	128.032	91.848	5201.868	2.261	102.800	5229.003
004649+2134	2	00 46 49.0	21 34 58.8	3.40	3.52	1.112	59.219	92.618	5170.682	...	...	...
011440+2708	2	01 14 40.3	27 08 06.0	3.47	3.49	3.261	144.113	109.525	3620.157	...	...	...
014206+1235	1	01 42 06.7	12 35 60.0	3.58	...	18.494	339.638	305.975	3065.943	...	...	...
014246+1309	1	01 42 46.8	13 09 18.0	3.65	...	8.125	152.775	129.193	807.862	...	...	...
014527+2531	2	01 45 27.6	25 31 19.2	3.54	3.57	7.417	110.014	91.536	3845.949	...	...	...
014729+2719	2	01 47 29.5	27 19 58.8	3.56	3.52	39.051	129.126	94.210	334.825	62.860	104.900	375.707
014847+1034	2	01 48 47.0	10 34 08.4	3.64	4.40	1.512	241.366	134.979	5245.624	1.587	146.500	5283.000
015011+2309	2	01 50 11.8	23 09 21.6	3.53	3.41	1.485	131.067	90.586	2863.507	2.934	87.690	2886.000
015105+1235	1	01 51 05.3	12 35 16.8	3.65	...	2.144	273.356	183.402	3280.678	...	...	...
015434+2312	2	01 54 34.3	23 12 14.4	3.51	3.49	...	...	...	...	3.482	95.810	5045.000
015443+1033	1	01 54 43.2	10 33 57.6	3.54	...	1.781	49.790	32.086	6086.814	...	...	...
015906+2523	2	01 59 06.2	25 23 09.6	3.72	3.50	3.205	247.220	231.736	5066.079	...	...	...
020022+2434	2	02 00 22.8	24 34 48.0	3.53	3.44	4.055	256.081	228.280	5134.388	...	...	...
020045+2809	2	02 00 45.2	28 09 57.6	3.43	3.46	2.069	66.911	48.819	5343.904	...	...	...
020148+2632	2	02 01 48.7	26 32 49.2	3.68	3.82	4.134	356.984	266.742	5021.338	...	...	...
020320+1837	2	02 03 20.2	18 37 48.0	5.80	3.51	...	...	...	...	10.250	86.490	2395.007
020320+2345	2	02 03 20.9	23 45 39.6	3.49	3.78	1.749	230.609	202.268	2792.723	3.095	115.600	2855.007
020405+2412	2	02 04 05.0	24 12 32.4	3.57	3.50	3.110	72.172	36.86	609.264	3.243	47.200	641.707
020918+2534	2	02 09 18.0	25 34 15.6	3.39	3.65	4.081	406.151	337.973	4942.253	...	...	...
022807+1935	2	02 28 07.7	19 35 34.8	3.47	3.55	12.519	415.978	394.993	4205.212	...	...	...
022859+2808	2	02 28 59.0	28 08 42.0	3.51	3.49	...	...	...	...	6.287	101.600	1036.000

Table 3. Optical Source Information

ID	Opt. cat. name	l	b	A_v	Diam ¹	Ell ¹	PA ¹
000330+2312	UGC 14	108.8479	-38.3574	0.406	102.675	0.392	-64.200
000407+2234	UGC 24	108.8361	-38.9910	0.287	60.010	0.408	-41.800
000623+2347	KUG 0003+235	109.8355	-37.9313	0.710	56.604	0.631	-50.000
000900+2348	NGC 9	110.5862	-38.0365	0.403	78.439	0.577	69.300
001622+2237	...	112.4226	-39.5372	0.379	42.926	0.583	9.100
002249+2310	...	114.4323	-39.2270	0.253	30.229	0.285	79.400
002526+2136	...	114.9329	-40.8636	0.293	23.963	0.339	53.300
003426+2436	UGC 337	118.0211	-38.1087	0.135	50.813	0.191	27.900
003751+0838	NGC 180	117.2043	-54.0786	0.240	126.776	0.320	81.400
003811+2523	...	119.1644	-37.3788	0.130	38.700	0.203	19.700
004649+2134	ESDO F540-04	121.5026	-41.2759	0.134	38.463	0.140	-78.700
011440+2708	ESDO F475-04	129.2796	-35.4502	0.247	53.476	0.114	55.400
014206+1235	NGC 658	141.7429	-48.4122	0.194	175.100	0.523	-64.300
014246+1309	UGC 1200 ³	141.7314	-47.8338	0.200	124.038	...	...
014527+2531	UGC 1230	137.9862	-35.7655	0.477	60.156	0.238	22.500
014729+2719	IC 1727	137.9526	-33.8985	0.340	333.036	0.691	22.450
014847+1034	UGC 1268	145.0964	-49.8188	0.517	86.773	0.673	-43.800
015011+2309	UGC 1294	140.0824	-37.7510	0.469	45.526	0.308	-33.600
015105+1235	UGC 1314	144.8569	-47.7283	0.303	80.921	0.587	72.200
015434+2312	LSBC F477-01	141.2767	-37.4084	0.469	34.931	0.221	84.600
015443+1033	ESDO F685-07	147.1993	-49.3008	0.491	51.559	0.237	-38.400
015906+2523	UGC 1462	141.6537	-35.0137	0.528	83.242	0.432	-13.000
020022+2434	IC 1764	142.3063	-35.6845	0.505	88.731	0.223	-71.300
020044+2809	...	141.0223	-32.2591	0.263	...	...	...
020148+2632	UGC 1510	141.9026	-33.7190	0.316	60.015	0.475	-40.100
020320+1837	UGC 1546	145.7422	-41.0354	0.325	70.216	0.057	-52.900
020320+2345	UGC 1538	143.4316	-36.2317	0.381	43.730	0.277	-25.600
020405+2412	UGC 1561	143.4377	-35.7533	0.362	118.698	0.294	10.007
020918+2534	UGC 1648	144.2080	-34.0635	0.325	90.083	0.454	-17.200
022807+1935	NGC 935 ³	152.1288	-37.7363	0.823	140.072	...	...
022859+2808	UGC 1958	147.7849	-29.9767	0.576	69.318	0.657	-72.400

Table 4. Sources Without an Identified Optical Counterpart

ID	b	A_v	Plate	Notes
020044+2809	-32.2591	0.263	POSS II	May be associated with large nearby galaxy
024328+2035	-35.1039	0.508	POSS II	Faint nebulosity behind bright star
025537+1938	-34.3980	6.438	POSS II	High A_v even though it is 34° out of the Plane
044753+2346	-13.5452	3.249	POSS I	High A_v
053017+2233	-6.3132	1.768	POSS II	Poor image due to bright star, but nothing obvious
055452+1509	-5.1948	1.379	POSS I	High stellar density, but nothing galaxy-like
055517+2526	0.0552	4.287	POSS II	Galactic Plane
060719+1936	-0.4254	6.776	POSS I	Galactic Plane
060733+0932	-5.2488	2.210	POSS I	Galactic Plane
061543+1110	-2.7069	3.419	POSS II	Galactic Plane
061729+2807	5.6839	2.313	POSS II	Galactic Plane
062054+2008	2.6385	3.416	POSS II	Galactic Plane
063549+1107	1.6357	6.332	POSS I	Galactic Plane
063603+1109	1.6983	7.149	POSS I	Galactic Plane
084411+2208	34.1423	0.166	POSS II	3 faint sources, unclear ID
084504+0932	29.5464	0.265	POSS II	Lots of small fuzzy things in field, unclear ID
090552+2520	39.8224	0.160	POSS II	Many fuzzy sources, unclear ID
142335+2131	68.3076	0.123	POSS II	2 faint sources straddling detection position
183229+2308	14.2863	0.513	POSS II	Possibly faint nebulosity behind bright star
184335+2007	10.6904	1.141	POSS II	High stellar density, but nothing galaxy-like
184829+1835	8.9901	1.379	POSS II	High stellar density, but nothing galaxy-like
192728+2012	1.5309	14.801	POSS I	Galactic Plane





This figure "figure2.jpg" is available in "jpg" format from:


<http://arXiv.org/ps/astro-ph/0004205v2>


This figure "figure3.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0004205v2>












This figure "figure7.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0004205v2>










This figure "figure10.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0004205v2>








This figure "figure13a.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0004205v2>

This figure "figure13b.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0004205v2>



This figure "figure15.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0004205v2>

